

ACUTE STROKE ADMISSION ORDERS

Admit to: Intermediate care unit () H61/floor ()

Diagnosis: _____ Condition: _____

Vitals: q _____ hrs. Neuro Checks: q _____ hrs.

Blood pressure: () Call H.O. for SBP > _____ SBP < _____ DBP > _____ DBP < _____

Activity: HOB < 30 deg () HOB < 60 deg () HOB > 60 deg ()

OOB to bathroom only () OOB to chair () Other _____

Allergies: _____

IV fluids: () Saline loc () IV normal saline @ _____ cc/hr with _____ mEq/L KCL

() IV 1/2 NS @ _____ cc/hr with _____ mEq/l KCL

() Other _____

Diet: () NPO () soft solids () pureed

() Thick liquids () no added Salt () NAS/low cholesterol

() Low cholesterol () CCDD (diabetic diet)

Medications: () IV heparin (25,000U/500 cc NS) @ _____ Units/hr

() SQ heparin 5000U BID (do not combine with IV heparin)

() Aspirin _____ mg PO/PR/png q day

() Coumadin _____ mg PO today

() Ticlid 250 mg PO BID

() Tylenol 650 mg PO/PR q 4 hr PRN

() Colace 100 mg PO q day PRN

() Pepcid 20 mg IV/PO BID (if NPO or critically ill)

Other medications: _____

Labs: () PTT _____ hrs after starting heparin () CXR if not done in last 6 months or symptomatic

() EKG () INR if on coumadin

Diagnostic Work Up: () CT head () Carotid ultrasounds () Transcranial dopplers (TCDs)

Preventive Therapy: () PAS stocking () Aspiration precautions

Rehab: () Consult physical therapy () Consult occupational therapy

() Consult speech therapy () Consult rehabilitation medicine

Discharge Planning: () Consult alternate site coordinator () Consult social services